

ASYMPTOTIC H -PLATEAU PROBLEM IN $\mathbb{H}^3$

BARIS COSKUNUZER

ABSTRACT. We show that for any Jordan curve Γ in $S_\infty^2(\mathbb{H}^3)$ with at least one smooth point, there exists an embedded H -plane $\mathcal{P}_H$ in $\mathbb{H}^3$ with $\partial_\infty \mathcal{P}_H = \Gamma$ for any $H \in [0, 1)$.

1. INTRODUCTION

There are two versions of the asymptotic Plateau problem. The first version asks the existence of a least area plane $\mathcal{P}$ in $\mathbb{H}^3$ asymptotic to a given simple closed Γ in $S_\infty^2(\mathbb{H}^3)$, i.e. $\partial_\infty \mathcal{P} = \Gamma$. In this version, there is a topological restriction on the surface $\mathcal{P}$ to be in the type of disk. The other version asks the existence of an area minimizing surface Σ in $\mathbb{H}^3$ asymptotic to a given collection of Jordan curves $\hat{\Gamma}$ in $S_\infty^2(\mathbb{H}^3)$, i.e. $\partial_\infty \Sigma = \hat{\Gamma}$. In the latter version, there is no a priori topological restriction on the surface Σ , hence Σ can have positive genus depending on given $\hat{\Gamma}$. Anderson gave positive answers to both questions 3 decades ago [A1, A2].

Constant mean curvature (CMC) surfaces are natural generalizations of minimal surfaces, and in many cases, the results related to minimal surfaces are studied to see whether they can be generalized to CMC setting. In our case, we will call this natural generalization as *asymptotic H -Plateau problem*. A decade after Anderson's result, the second version of the asymptotic Plateau problem was generalized to CMC case by Tonegawa [To]. Tonegawa showed that for any given collection of Jordan curves $\hat{\Gamma}$ in $S_\infty^2(\mathbb{H}^3)$, there exists a minimizing H -surface Σ_H in $\mathbb{H}^3$ with $\partial_\infty \Sigma_H = \hat{\Gamma}$ where $H \in [0, 1)$. Indeed, both Anderson and Tonegawa used geometric measure theory methods, and the solutions are automatically smoothly embedded surfaces by the regularity results of GMT. The survey [C2] gives a fairly complete account of the old and new results on the problem.

On the other hand, for the generalization of the first (plane) version to the CMC case, the only result came out a few years ago by Cuscheri [Cu]. He showed the existence of *immersed H -planes* asymptotic to given smooth Jordan curve in $S_\infty^2(\mathbb{H}^3)$ by using PDE techniques.

The author is supported by Fulbright Grant, and TUBITAK 2219 Grant.

In this paper, we give positive answer to the asymptotic H -Plateau problem for a larger family of curves. Furthermore, we show that these solutions are indeed *embedded*.

Theorem 1.1. *Let Γ be a simple closed curve Γ in $S_\infty^2(\mathbb{H}^3)$ with at least one C^1 -smooth point. Then, for any $H \in [0, 1)$, there exists a properly embedded H -plane $\mathcal{P}_H$ in $\mathbb{H}^3$ with $\partial_\infty \mathcal{P}_H = \Gamma$.*

Our techniques are also valid for $H = 0$ case, and we are able to reprove the existence of least area planes in $\mathbb{H}^3$. Hence, with this result, we filled the gap in Anderson's result in [A2], too (See Remark 3.1).

On the other hand, our proof is indeed for $-1 < H < 1$ by considering the orientation. Hence, if you forget the sign of the mean curvature, the above theorem shows that for a given Jordan curve Γ in $S_\infty^2(\mathbb{H}^3)$ and $H \in (0, 1)$, there exist a pair of complete, embedded H -planes $\mathcal{P}_H^+$ and $\mathcal{P}_H^-$ with $\partial_\infty \mathcal{P}_H^\pm = \Gamma$ (See Remark 4.3).

Recently, Meeks, Tinaglia and the author constructed nonproperly embedded H -planes in $\mathbb{H}^3$ for any $H \in [0, 1)$ [CMT], where the asymptotic boundary is a pair of infinite lines in $S_\infty^2(\mathbb{H}^3)$. Here, we also show that if the Jordan curve Γ in $S_\infty^2(\mathbb{H}^3)$ is smooth enough, then the minimizing H -planes $\mathcal{P}_H$ with $\partial_\infty \mathcal{P}_H = \Gamma$ are *properly embedded* in $\mathbb{H}^3$ (Corollary 4.5).

In the final section, we discuss the questions of the *generic uniqueness* of the H -planes in $\mathbb{H}^3$, and the *foliations of $\mathbb{H}^3$* with H -planes, and give an outline to solve them.

The organization of the paper is as follows: In the next section, we go over the basic notions, and the related results. In Section 3, we prove the main theorem for least area planes ($H = 0$ case). In Section 4, we show the existence of embedded H -planes in $\mathbb{H}^3$. Finally in Section 5, we give some concluding remarks.

2. PRELIMINARIES

In this section, we will go over the related results which will be used in the following sections. For further details of the notions, and the results used in this paper, one can see the survey [C2].

Definition 2.1. Let M be a 3-manifold.

- *Minimal surface:* A surface S in M is minimal if the mean curvature vanishes everywhere on S .
- *Least Area Disk:* A compact disk D in M with $\partial D = \Gamma$ is the least area disk in M if it has the smallest area among the disks in M with the boundary Γ .

Note that minimal surfaces are the critical points of the area functional. Least area disks and area minimizing surfaces are the minimum of the area functional in the corresponding spaces.

A natural generalization of minimal surfaces are CMC surfaces (H -surfaces). They can be defined as the critical points of the area functional with a volume constraint as follows. For an immersion $u : D^2 \rightarrow M$, the critical points of the following variational problem are immersed disks with constant mean curvature H [Gu].

$$\mathcal{F}_H(u) = \int_{D^2} (|u_x|^2 + |u_y|^2) + \frac{4}{3}H[u \cdot (u_x \times u_y)] dx dy$$

Here, the second summand in the integral represents the volume constraint.

We can reformulate this variational problem in a different way so that it will be independent of the parametrization of the surface [C5]. Let Σ be a surface in M with boundary α . We fix a surface T in M with $\partial T = \alpha$, and define Ω to be the domain bounded by T and Σ . Again, let

$$\mathcal{I}_H(\Sigma) = \text{Area}(\Sigma) + 2H \text{Vol}(\Omega)$$

If Σ is a critical point of the functional $\mathcal{I}_H$ for any variation f , then this will imply Σ has constant mean curvature H . Note also that critical point of the functional $\mathcal{I}_H$ is independent of the choice of the surface T since if $\hat{\mathcal{I}}_H$ is the functional which is defined with a different surface $\hat{T}$, then $\mathcal{I}_H - \hat{\mathcal{I}}_H = C$ for some constant C . Note that to keep the solution surface away from T , one needs a convexity condition on T (e.g. H_0 -convex for $H_0 > H$) to employ the maximum principle [C5].

Definition 2.2. Let M be a 3-manifold.

- *H -surface:* A surface S in M is H -surface if the mean curvature is equal to H everywhere on S .
- *Minimizing H -disk:* A compact disk D in M with $\partial D = \Gamma$ is a minimizing H -disk in M if $\mathcal{I}_H(D)$ (or equivalently $\mathcal{F}_H(D)$) has the smallest value among the disks in M with the boundary Γ .

2.1. Embedded solutions to the H -Plateau problem: Here, we will quote the generalization of Meeks-Yau's embeddedness result [MY] to H -disks.

Definition 2.3 (H_0 -convex domains). Let Ω be a compact 3-manifold with piecewise smooth boundary. We call Ω an H_0 -convex domain if

- The mean curvature vector $\mathbf{H}$ always points towards inside Ω along the smooth parts of $\partial\Omega$,
- The mean curvature $|\mathbf{H}(p)| \geq H_0$ for any smooth point $p \in \partial\Omega$,
- Along the nonsmooth parts of $\partial\Omega$, the inner dihedral angle is less than π .

The following lemma is on the embeddedness of the solutions of H -Plateau problem for H -extreme curves.

Lemma 2.4. [C5] *Let M be a compact H_0 -convex ball. Let Γ be a Jordan curve in ∂M . Then, for any $H \in [0, H_0)$, there exists a minimizing H -disk Σ_H in M with $\partial\Sigma_H = \Gamma$, and any such Σ_H is embedded.*

The following lemma is known as maximum principle.

Lemma 2.5. [Gu] *[Maximum Principle] Let Σ_1 and Σ_2 be two surfaces in a Riemannian 3-manifold which intersect at a common point tangentially. If Σ_2 lies in positive side (mean curvature vector direction) of Σ_1 around the common point, then H_1 is strictly less than H_2 ($H_1 < H_2$) where H_i is the mean curvature of Σ_i at the common point.*

2.2. H -Planes in $\mathbb{H}^3$: Now, we restrict ourselves to $\mathbb{H}^3$. Let Γ be a simple closed curve in $S_\infty^2(\mathbb{H}^3)$. Γ separates $S_\infty^2(\mathbb{H}^3)$ into two open disks, say D^+ and D^- . We fix an orientation, and consider the mean curvature H with sign depending on the direction of the mean curvature vector, i.e. $-1 < H < 1$ such that $\mathcal{P}_H \sim D^- \Rightarrow H \sim +1$ and $\mathcal{P}_H \sim D^+ \Rightarrow H \sim -1$ [C3, To].

Now, by using the definitions above, we define the least area planes and the minimizing H -planes in $\mathbb{H}^3$.

Definition 2.6 (Least Area Plane). Let $\mathcal{P}$ be a complete surface in $\mathbb{H}^3$ which is topologically in the type of a disk. We call $\mathcal{P}$ a least area plane in $\mathbb{H}^3$, if any compact subdisk D in $\mathcal{P}$ is a least area disk.

Definition 2.7 (Minimizing H -plane). Fix $H \in (-1, 1)$. Let $\mathcal{P}_H$ be a complete surface in $\mathbb{H}^3$ which is topologically in the type of a disk. We call $\mathcal{P}_H$ a minimizing H -plane in $\mathbb{H}^3$, if any compact subdisk D in $\mathcal{P}_H$ is a minimizing H -disk.

For a given surface S in $\mathbb{H}^3$, we define the *asymptotic boundary* of S as follows. If $\overline{\mathbb{H}^3} = \mathbb{H}^3 \cup S_\infty^2(\mathbb{H}^3)$ is the natural (geodesic) compactification of $\mathbb{H}^3$, and $\overline{S}$ is the closure of S in $\overline{\mathbb{H}^3}$, then the asymptotic boundary $\partial_\infty S$ of S defined as $\partial_\infty S = \overline{S} \cap S_\infty^2(\mathbb{H}^3)$.

Now, we define the shifted convex hulls $CH_H(\Gamma)$ as generalizations of the convex hulls in $\mathbb{H}^3$ [C2, C3]. Fix $H \in (-1, 1)$. Let Γ be a Jordan curve in $S_\infty^2(\mathbb{H}^3)$. Let α be a round circle in $S_\infty^2(\mathbb{H}^3)$ with $\alpha \cap \Gamma = \emptyset$. Let $\mathcal{P}_H^\alpha$ be the unique H -plane in $\mathbb{H}^3$ with $\partial_\infty \mathcal{P}_H^\alpha = \alpha$. α separates $S_\infty^2(\mathbb{H}^3)$ into two open disks Δ_α^+ and Δ_α^- . Similarly, $\mathcal{P}_H^\alpha$ divides $\mathbb{H}^3$ into two domains $\Omega_H^{\alpha+}$ and $\Omega_H^{\alpha-}$ where $\partial_\infty \Omega_H^{\alpha\pm} = \Delta_\alpha^\pm$. We will call these regions as *H -shifted halfspaces*. If $\Gamma \subset \Delta_\alpha^+$, then we will call $\Omega_H^{\alpha+}$ a *supporting H -shifted halfspace*. Similarly, if $\Gamma \subset \Delta_\alpha^-$, then we will call $\Omega_H^{\alpha-}$ a *supporting H -shifted halfspace*.

Definition 2.8 (Shifted Convex Hull). Let Γ be a simple closed curve in $S_\infty^2(\mathbb{H}^3)$. Fix $H \in (-1, 1)$. Then the H -shifted convex hull of Γ , $CH_H(\Gamma)$ is defined as the intersection of all supporting closed H -shifted halfspaces $\Omega_H^{\alpha\pm}$ of $\mathbb{H}^3$. For $H = 0$, this is the usual convex hull definition in $\mathbb{H}^3$, i.e. $CH(\Gamma) = CH_0(\Gamma)$.

Now, the generalization of convex hull property of minimal surfaces in $\mathbb{H}^3$ to H -surfaces in $\mathbb{H}^3$ is as follows [C3, To].

Lemma 2.9. [To], [C3] *Let Σ be a H -surface in $\mathbb{H}^3$ where $\partial_\infty \Sigma = \Gamma$ and $|H| < 1$. Then Σ is in the H -shifted convex hull of Γ , i.e. $\Sigma \subset CH_H(\Gamma)$.*

Remark 2.10. Notice that the result is true for any H -surface. This is a straightforward generalization of the convex hull property for minimal surfaces. In particular, if Σ is an H -surface in $\mathbb{H}^3$ with $\partial_\infty \Sigma_H = \Gamma$, then Σ cannot go into a nonsupporting H -shifted halfspaces of Γ , as we can foliate these halfspaces with H -planes, and the first point of touch gives a contradiction with the maximum principle.

3. EXISTENCE OF LEAST AREA PLANES

In this section, we will focus on $H = 0$ case. In other words, we will consider the original asymptotic Plateau problem, and show the existence of smoothly embedded least area planes $\mathcal{P}$ in $\mathbb{H}^3$ with $\partial_\infty \mathcal{P} = \Gamma$ for a given Jordan curve Γ in $S_\infty^2(\mathbb{H}^3)$.

Note that Gabai showed the existence of least area planes in $\mathbb{H}^3$ in [Ga] by using Hass and Scott's techniques. Recently, Ripoll and Tomi also showed the existence of complete embedded minimal planes in Hadamard manifolds [RT1].

In this paper, we will adapt Anderson's techniques in [A2] to construct minimizing H -planes. To generalize his techniques, we need to fill a gap in the proof for least area plane case. Following remark explains the problem.

Remark 3.1. (Gap in [A2, Theorem 4.1]) Anderson showed the existence of least area planes in [A2, Theorem 4.1]. He basically generalized the techniques he used for absolutely area minimizing surface case to the plane case. In particular, let Γ be a Jordan curve in $S_\infty^2(\mathbb{H}^3)$, and $\{D_n\}$ be a sequence of least area disks in $B_n(0)$ with $\partial D_n = \gamma_n \subset \partial B_n(0)$ where $\gamma_n \rightarrow \Gamma$. Then, the idea is to show the existence of a subsequence of $\{D_n\}$ converging to a least area plane $\mathcal{P}$ in $\mathbb{H}^3$ with $\partial_\infty \mathcal{P} = \Gamma$. In particular, for every fixed compact domain K in $\mathbb{H}^3$, he showed the sequence $D_n \cap K = D_n^K$ has a subsequence converging to a smooth disk in K by using the compactness and regularity results of GMT. Then, by using the diagonal sequence argument, he obtained a limit least area plane $\mathcal{P}$ with $\partial \mathcal{P} = \Gamma$.

However, to use the compactness result in this approach, one needs a uniform area bound on the disks $\{D_n^K\}$. Let Γ_n^K be the collection of simple closed curves $D_n \cap \partial K$. If $\{\Sigma_n\}$ was a sequence of area minimizing surfaces, then the area of ∂K would be a uniform area bound for $\{\Sigma_n^K\}$ as Γ_n^K bounds a surface in ∂K and Σ_n^K is absolutely area minimizing surface with boundary Γ_n^K . Hence, for any n , $Area(\Sigma_n^K) \leq Area(\partial K)$. However, in the disk case $D_n \cap K$ may contain many disks or planar surfaces in K , and the area of ∂K cannot give an upper bound for $Area(D_n^K)$. In particular, the estimate (4.2) ($M(D_i \setminus B_r) \leq \frac{1}{2}Area(S(r))$) in [A2, Theorem 4.1] is not valid in general.

For example, if D_n^K is $2k$ disjoint disks close to the equator disk in $K = B_R$, and the area of D_n^K would be close to the area of k equator disks, which is much larger than the area of ∂B_R . The main difference with the area minimizing case is that if we have k annuli $A_1, \dots, A_k$ in ∂K bounding Γ_n^K , we cannot compare the sum of the areas of the disks with the sum of the areas of the annuli. Because if we replace two disks with an annulus in D_n we would get a genus 1 surface, which is no longer a disk. So, the area of an annulus cannot be compared with the area of the two disks, because of the restriction of the topology on $\{D_n\}$. Since there is no restriction on the topology of the surface for area minimizing surfaces $Area(\partial K)$ gives a uniform bound, but in the least area disk case, $Area(\partial K)$ does not give a uniform bound for $\{D_n^K\}$.

Now, we will show the existence of least area plane in $\mathbb{H}^3$ by using Anderson's techniques. In order to get the uniform area bound on $\{D_n^K\}$ for fixed compact set K , we will use the ideas in [C1].

Theorem 3.2. *Let Γ be a simple closed curve with at least one smooth (C^1) point in $S_\infty^2(\mathbb{H}^3)$. Then, there exists a properly embedded least area plane $\mathcal{P}$ in $\mathbb{H}^3$ with $\partial_\infty \mathcal{P} = \Gamma$.*

Proof: We will define a special sequence of least area disks $\{D_n\}$ where their restriction to a compact subset K has a uniform area bound. Then, by following [A2], we get a least area plane.

Notation and the Setting: Let Γ be a Jordan curve with at least one smooth (C^1) point in $S_\infty^2(\mathbb{H}^3)$. Let $CH(\Gamma)$ be the convex hull of Γ in $\mathbb{H}^3$. Near the smooth point $p \in \Gamma$, we can have two sufficiently small and close round circles τ^+ and τ^- in the opposite sides of Γ such that $\tau^+ \cup \tau^-$ bounds a least area annulus $\mathcal{A}$ in $\mathbb{H}^3$ [Wa]. Hence, $\mathcal{A}$ goes through $CH(\Gamma)$.

Fix a point O in $CH(\Gamma)$. Let B_n be the closed ball in $\mathbb{H}^3$ of radius n with center O . Then, for sufficiently large N_0 , $\mathcal{A} \cap CH(\Gamma)$ is in B_{N_0} . Let $\partial CH(\Gamma) = \partial^+ CH(\Gamma) \cup \partial^- CH(\Gamma)$. For $n > N_0$, let $\alpha_n^+ = \partial B_n \cap \partial^+ CH(\Gamma)$ and $\alpha_n^- = \partial B_n \cap \partial^- CH(\Gamma)$. Then, since $\mathcal{A}$ is a least area

annulus, by [MY], the pair $\alpha_n^+ \cup \alpha_n^-$ bounds a least area annulus $\mathcal{A}_n$ in $CH(\Gamma)$.

Let Ω_n be the compact region which $\mathcal{A}_n$ separates from $CH(\Gamma)$. Let γ_n be an essential, smooth, simple closed curve in $\mathcal{A}_n$. Then by construction, $\gamma_n \rightarrow \Gamma$ as $\partial_\infty CH(\Gamma) = \Gamma$. Let D_n be the least area disk in Ω_n with $\partial D_n = \gamma_n$ [MY]. By construction, D_n is also a least area disk in $\mathbb{H}^3$.

Now, we claim that there exists a subsequence of $\{D_n\}$ which converges to a least area plane $\mathcal{P}$ in $\mathbb{H}^3$ with $\partial_\infty \mathcal{P} = \Gamma$. We will follow the proof of [A2, Theorem 4.1]. Hence, if we show the estimate (4.2) [A2, Theorem 4.1] ($M(D_n \sqcup \mathbf{B}_r) \leq \frac{1}{2} \text{Area}(\partial \mathbf{B}_r)$) is valid for our sequence, we are done.

Now, let β be a transversal arc in $CH(\Gamma)$ connecting $\partial^+ CH(\Gamma)$ and $\partial^- CH(\Gamma)$ through the point O (centers of the balls $\mathbf{B}_n$). Let l be the length of β . Consider the following lemma from [C1].

Let D_r be a least disk in $\mathbf{B}_r$ with $\partial D_r \subset \partial \mathbf{B}_r \cap CH(\Gamma)$. Then, we call D_r *nonseparating with respect to Γ in $\mathbf{B}_r$* (say wrt- Γ), if ∂D_r is not an essential curve in the annulus $\partial \mathbf{B}_r \cap CH(\Gamma)$.

Lemma 3.3. [C1, Lemma 4.1] *Let Γ be a Jordan curve in $S_\infty^2(\mathbb{H}^3)$ with at least one smooth (C^1) point. Let D_r be a least area disk, nonseparating wrt- Γ in $\mathbf{B}_r$ for $r > N_0$. Then there is a monotone increasing function $F : [N_0, \infty) \rightarrow \mathbb{R}^+$ such that $F(r) \rightarrow \infty$ as $r \rightarrow \infty$, and $d(O, D_r) > F(r)$ where d is the distance.*

Let $R_0 > 0$ be sufficiently large so that $F(R_0) > l$. Now, we will prove the uniform bound $|D_n \cap \mathbf{B}_r| < C_r$ for $r > R_0$ where $|\cdot|$ represents the area.

By [A2, Lemma 4.2], for a given least area disk D_n , $D_n \cap \mathbf{B}_r$ is a collection of disjoint embedded disks for any generic $r > 0$. This is simply because $D_n \cap \mathbf{B}_r$ is a surface for generic r , and by the convexity of $\mathbf{B}_r$, any component in $D_n \cap \mathbf{B}_r$ must be a disk.

We claim that for $n > \max\{N_0, R_0\}$, $D_n \cap \mathbf{B}_r$ is just a disk (only one component) for any generic $r > R_0$. Assume on the contrary, and let E_1, E_2 be two such disks in $D_n \cap \mathbf{B}_{r_0}$. Since D_n is an embedded disk in $\mathbf{B}_n$, there is a path α connecting E_1 and E_2 in D_n . Let $r_0 < r' < n$ be the smallest radius such that E_1 and E_2 are in the same component $\hat{E}$ of $D_n \cap \mathbf{B}_{r'}$. Hence, $\hat{E}$ is a least area disk, nonseparating wrt- Γ in $\mathbf{B}_{r'+\epsilon}$ for sufficiently small $\epsilon > 0$. Hence by the Lemma 3.3, $d(O, \hat{E}) > F(r')$. Hence, we get $d(O, \hat{E}) \leq d(O, E_i) \leq l < F(R) < F(r')$ which gives a contradiction.

This proves that for $n > \max\{N_0, R_0\}$ and for any $r > R_0$, $D_n \cap \mathbf{B}_r$ is a disk (only one component). Hence, for any fixed $r > R_0$, $|D_n \cap \mathbf{B}_r| \leq |\partial \mathbf{B}_r| = C_r$, which gives the desired uniform bound. This proves the estimate (4.2) of [A2, Theorem 4.1] is valid for our sequence $\{D_n\}$. The proof follows. $\square$

4. MINIMIZING H -PLANES

In this section, we will show the existence of the solutions of asymptotic H -Plateau problem in $\mathbb{H}^3$ for $H \in (-1, 1)$. In particular, we will generalize the techniques in the previous section to the CMC case, and show that for any simple closed curve Γ in $S_\infty^2(\mathbb{H}^3)$ with one smooth point, there exists an embedded H -plane $\mathcal{P}_H$ with $\partial_\infty \mathcal{P}_H = \Gamma$. First, we need to generalize Lemma 3.3 proven in [C1] to the minimizing H -planes in $\mathbb{H}^3$.

We will use the same notation, i.e. let $O, \mathbf{B}_r, \mathcal{A}_r, N_0$ be as in the previous section. Fix $H \in (-1, 1)$. Let $\hat{A}_\Gamma^H = \partial \mathbf{B}_r \cap CH_H(\Gamma)$ be an annulus in $\partial \mathbf{B}_r$. Again, let D_r be a minimizing H -disk in $\mathbf{B}_r$ with $\partial D_r \subset \hat{A}_\Gamma^H$. Then, we call D_r *nonseparating wrt- Γ in $\mathbf{B}_r$* , if ∂D_r is not an essential curve in the annulus $\hat{A}_\Gamma^H$.

Lemma 4.1. *Fix $H \in (-1, 1)$. Let Γ be a Jordan curve with at least one smooth (C^1) point in $S_\infty^2(\mathbb{H}^3)$. For $r > N_0$, let D_r be a minimizing H -disk, and nonseparating wrt- Γ in $\mathbf{B}_r$. Then there is a monotone increasing function $F : [N_0, \infty) \rightarrow \mathbb{R}^+$ such that $F(r) \rightarrow \infty$ as $r \rightarrow \infty$, and $d(O, D_r) > F(r)$ where d is the distance.*

Proof: We will adapt the proof of Lemma 3.3 to this case. Lemma 3.3 finishes the $H = 0$ case. Hence, we can take $H > 0$. For $H < 0$, the same proof works by changing the orientation. Fix $H > 0$.

Recall that $\mathcal{A}_r$ is the least area annulus in $\mathbf{B}_r$ with $\partial \mathcal{A}_r = \alpha_r^+ \cup \alpha_r^-$ for generic $r > N$. Let $\hat{A}_r$ be the annulus in $\partial \mathbf{B}_r$ with $\partial \hat{A}_r = \alpha_r^+ \cup \alpha_r^-$. Let $F(r) = d(O, \mathcal{A}_r)$.

We claim that for any minimizing H -disk D_r which is nonseparating wrt- Γ in $\mathbf{B}_r$, $D_r \cap \mathcal{A}_r = \emptyset$, i.e. D_r stays in the solid torus $\mathcal{U}_r$ in $\mathbf{B}_r$ with $\partial \mathcal{U}_r = \mathcal{A}_r \cup \hat{A}_r$; See Figure 1. In particular, this shows that $d(O, D_r) > F(r)$, and the proof will follow.

Now, let D_r be a minimizing H -disk and nonseparating wrt- Γ in $\mathbf{B}_r$ with $\partial D_r = \gamma_r \subset \hat{A}_r$. Since D_r is nonseparating wrt- Γ , γ_r is a not an essential curve in $\hat{A}_r$. In other words, γ_r bounds a disk E_r in $\hat{A}_r$; see Figure 1.

Let Δ_r be the region in $\mathbf{B}_r$ with $\partial \Delta_r = D_r \cup E_r$. Since D_r is a minimizing H -disk in $\mathbf{B}_r$ with $H > 0$, $I_H(D_r) = |D_r| + 2H|\Delta_r|$ is the smallest among all the disks in $\mathbf{B}_r$ with boundary γ_r . Here, $|\cdot|$ represents the area or the volume of the corresponding region.

Assume that $d(O, D_r) < d(O, \mathcal{A}_r)$. Recall that $\mathcal{U}_r$ is the region in $\mathbf{B}_r$ with $\partial \mathcal{U}_r = \mathcal{A}_r \cup \hat{A}_r$. Let $\Delta'_r = \Delta_r \cap \mathcal{U}_r$. Then as $d(O, D_r) < d(O, \mathcal{A}_r)$, $|\Delta'_r| < |\Delta_r|$. Furthermore, let $T' = \mathcal{A}_r \cap \partial \Delta'_r$ be the planar region in $\mathcal{A}_r$ with $\partial T' = \beta$. Let T be the planar region in D_r with $\partial T = \beta$. Since D_r is a nonseparating disk, and $\mathcal{A}_r$ is an annulus, $D' = (D_r - T) \cup T'$ is also a disk in $\mathbf{B}_r$ with $\partial D' = \gamma_r$. Furthermore, as $\mathcal{A}_r$ is a least area annulus,




FIGURE 1. If γ_r is a nonessential curve in $\hat{A}_r$, the nonseparating minimizing H -disk D_r in B_r with $\partial D_r = \gamma_r$ must belong to $\mathcal{U}_r$. This shows that for any such D_r in B_r , $d(O, D_r) > d(O, \mathcal{A}_r) = F(r)$.

$|T'| < |T|$ and hence $|D'| < |D_r|$. This implies $\mathcal{I}_H(D') = |D'| + 2H|\Delta'_r| < |D_r| + 2H|\Delta'_r| = \mathcal{I}_H(D_r)$. However, D_r is a minimizing H -disk in B_r . This is a contradiction. This proves $D_r \cap \mathcal{A}_r = \emptyset$, and $D_r \subset \mathcal{U}_r$. Hence, this shows that $d(O, D_r) > d(O, \mathcal{A}_r) = F(r)$. The lemma follows. $\square$

Now, we prove the main result of the paper.

Theorem 4.2. *[Existence of H -Planes] Let Γ be a Jordan curve in $S^2_\infty(\mathbb{H}^3)$ with at least one smooth (C^1) point. Let $H \in (-1, 1)$. Then, there exists a properly embedded minimizing H -plane $\mathcal{P}_H$ in $\mathbb{H}^3$ with $\partial_\infty \mathcal{P}_H = \Gamma$.*

Proof: Fix $H \in (-1, 1)$. Given Γ in $S^2_\infty(\mathbb{H}^3)$, we will use the same setup as before, i.e. let $\mathcal{A}, \mathcal{A}_r, \hat{A}_r, O, B_r, N_0$ be as in the previous lemma.

Let $\Omega_n = B_n \cap CH_H(\Gamma)$ for $n > N_0$. Let γ_n be an essential smooth curve in $\hat{A}_n \subset \partial\Omega_n$. Then by construction, $\gamma_n \rightarrow \Gamma$ as $\partial_\infty CH_H(\Gamma) = \Gamma$. Recall that the mean curvature of the geodesic sphere of radius R is $\coth R$. Hence, B_n is 1-convex for any n . Then, by Lemma 2.4, there exists an embedded minimizing H -disk D_n in B_n with $\partial D_n = \gamma_n$. Here, the sign of H determines the minimizing H -disk as there are two minimizing H -disks in B_n facing each other for $H \neq 0$ [C5]. Furthermore, since $\gamma_n \subset CH_H(\Gamma)$, by Lemma 2.9 (see also Remark 2.10), $D_n \subset CH_H(\Gamma)$. Hence, this implies $D_n \subset \Omega_n = B_n \cap CH_H(\Gamma)$.

We claim that the sequence of embedded minimizing H -disks $\{D_n\}$ converges to a minimizing H -plane $\mathcal{P}_H$. First, we claim that the sequence of minimizing H -disks $\{D_n\}$ has a convergent subsequence in any compact

set K in $\mathbb{H}^3$. Then, by using the diagonal sequence argument as before, we will get a limit minimizing H -plane in $\mathbb{H}^3$.

Consider the closed balls $\mathbf{B}_K$ with center O . By using convexity of $\mathbf{B}_K$, for generic $K > N_0$, we can assume $D_n \cap \mathbf{B}_K$ is a collection of disks for any n by [A2, Lemma 4.2].

Now, let β be a transversal arc in $CH_H(\Gamma)$ connecting $\partial^+ CH_H(\Gamma)$ and $\partial^- CH_H(\Gamma)$ through the point O . Let l be the length of β . Fix a generic $K_0 > N_0$ such that $F(K_0) > l$. Then by Lemma 4.1, and the proof of Theorem 3.2, $D_n^0 = D_n \cap \mathbf{B}_{K_0}$ is a closed disk (only one component) for any n . Hence, the sequence of minimizing H -disks $\{D_n^0\}$ has a uniform area bound, say $|D_n^0| < |\partial \mathbf{B}_{K_0}|$. Then, by following the proof of [A2, Theorem 4.1], the existence of smoothly embedded minimizing H -plane Σ_H can be shown as follows:

With the uniform area bound, the sequence of minimizing H -disks $\{D_n^0\}$ has a convergent subsequence in B_{K_0} by the compactness theorem for integral currents. Hence by considering them as integral currents, a subsequence of $\{D_n^0\}$ converges to a properly embedded minimizing H -disk D^0 in $\mathbf{B}_{K_0}$. Let K_i be a monotone increasing sequence with $K_i \nearrow \infty$. For $K_1 > K_0$, by starting with this subsequence, get another subsequence converging on $\mathbf{B}_{K_1}$. By iterating this process and diagonal sequence argument, we get a sequence of integral currents $\{D_n\}$ converges on compacts to the integral current Σ_H in $\mathbb{H}^3$. By Allard's regularity, the convergence is smooth on compact sets. Also, the asymptotic boundary of the support of Σ_H is Γ by the convex hull property (Lemma 2.9), i.e. $\partial_\infty \Sigma_H = \Gamma$ as $\partial_\infty CH_H(\Gamma) = \Gamma$.

The limit of minimizing H -disks Σ_H is a minimizing H -surface. Hence, by [Gu], for any point p in the support of Σ_H , there exists $\epsilon > 0$ with $\mathbf{B}_\epsilon(p) \cap \Sigma_H$ is a smooth embedded disk. Hence, the support of Σ_H is smoothly embedded surface. Finally, since the convergence is smooth in compact sets, and $\{D_n\}$ is a sequence of embedded disks, Σ_H is a complete minimizing H -plane in $\mathbb{H}^3$ with $\partial_\infty \Sigma_H = \Gamma$. The proof follows. $\square$

Remark 4.3. [Pair of H -planes] Notice that if you forget the sign of the mean curvature H , the theorem above shows that for a given Jordan curve Γ in $S_\infty^2(\mathbb{H}^3)$, there exist two minimizing H -planes $\mathcal{P}_H^+$ and $\mathcal{P}_H^-$ with $\mathcal{P}_H^\pm = \Gamma$ for $H \in (0, 1)$. Furthermore, these H -planes are disjoint, and the convex sides are facing each other.

Remark 4.4. The above result was shown for special families of curves in $S_\infty^2(\mathbb{H}^3)$ like star-shaped curves [GS], "mean convex" curves [NS], and graph over a line [RT2], where they showed that the area minimizing surface Σ_H is indeed a graph over a geodesic subspace in $\mathbb{H}^3$.

Corollary 4.5. *[Properly Embeddedness] Let Γ be a $C^{3,\alpha}$ smooth Jordan curve in $S_\infty^2(\mathbb{H}^3)$. Let $\mathcal{P}_H$ be a minimizing H -plane in $\mathbb{H}^3$ with $\partial_\infty \mathcal{P}_H = \Gamma$. Then, $\mathcal{P}_H$ is properly embedded in $\mathbb{H}^3$.*

Proof: By [To], since Γ is $C^{3,\alpha}$ smooth, $\mathcal{P}_H$ is regular near infinity. In particular, there exists a $\rho > 0$ such that in the upper half space model of $\mathbb{H}^3$, $\mathcal{P}_H \cap \{z < \rho\}$ is a graph over $\Gamma \times (0, \rho)$. Then, for sufficiently large N , $\partial \mathbf{B}_N \cap \mathcal{P}_H$ is a Jordan curve γ in $\partial \mathbf{B}_N$. Hence, $\mathbf{B}_N \cap \mathcal{P}_H = \mathcal{D}$ is a minimizing H -disk in $\mathbf{B}_N$ with $\partial \mathcal{D} = \gamma$ by the definition of minimizing H -plane. By Lemma 2.4, $\mathcal{D}$ is properly embedded. Since $\mathcal{P}_H - \mathcal{D}$ is a graph over $\Gamma \times (0, \rho)$ in the upper half space model [To], the proof follows. $\square$

Remark 4.6. Indeed, it can be showed that the above result is true for far more generality. By using the techniques in [C1], one can naturally generalize the above result to Jordan curves in $S_\infty^2(\mathbb{H}^3)$ with at least one smooth (C^1) point. Even though there is no regularity near infinity in that case, by using Lemma 4.1, the arguments in [C1] can easily be adapted.

Note also that, Meeks, Tinaglia and the author recently showed that there exists a *nonproperly* embedded complete H -plane Σ_H in $\mathbb{H}^3$ for any $H \in [0, 1)$ [CMT]. In particular, Σ_H is an H -plane between two rotationally invariant H -catenoids $\mathcal{C}_1$ and $\mathcal{C}_2$ where Σ_H spirals into $\mathcal{C}_1$ in one end, and spirals into $\mathcal{C}_2$ in the other end. Hence, $\partial_\infty \Sigma_H$ is a pair of infinite lines l^+ and l^- in $S_\infty^2(\mathbb{H}^3)$. Here, if $\partial_\infty \mathcal{C}_i = \alpha_i^+ \cup \alpha_i^-$, and $A^\pm$ is the annuli in $S_\infty^2(\mathbb{H}^3)$ with $\partial A_i^\pm = \alpha_1^\pm \cup \alpha_2^\pm$, then $l^+ \subset A^+$ and $l^- \subset A^-$ where $l^\pm$ spirals into $\alpha_1^\pm$ in one end, and spirals into $\alpha_2^\pm$ in the other end.

5. FINAL REMARKS

5.1. Generic Uniqueness of minimizing H -Planes.

The generic uniqueness results for minimizing H -surfaces in $\mathbb{H}^3$ [C3] can naturally be generalized to our context, i.e minimizing H -planes. In particular, for fixed $H \in (-1, 1)$, let Σ_1 and Σ_2 be minimizing H -planes with $\partial_\infty \Sigma_i = \Gamma_i$ where Γ_1 and Γ_2 are disjoint simple closed curves in $S_\infty^2(\mathbb{H}^3)$. Then, by using Meeks-Yau exchange roundoff trick, it can be showed Σ_1 and Σ_2 are disjoint, too. By using this, and similar ideas to [C3], it can be showed that any simple closed curve Γ bounds either a unique minimizing H -plane Σ in $\mathbb{H}^3$, or there are two canonical disjoint minimizing H -planes Σ^+ and Σ^- with $\partial_\infty \Sigma^\pm = \Gamma$. Hence, foliating an annular neighborhood of Γ in $S_\infty^2(\mathbb{H}^3)$ with simple closed curves, and considering the canonical H -planes constructed, one can get generic uniqueness result as in [C3]. In particular, this shows that *for fixed $H \in (-1, 1)$, a generic Jordan curve Γ in $S_\infty^2(\mathbb{H}^3)$ bounds a unique minimizing H -plane Σ in $\mathbb{H}^3$ with $\partial_\infty \Sigma = \Gamma$.*

5.2. Foliations of $\mathbb{H}^3$ by H -planes.

Similar to the previous part, it is also possible to show that two H -planes in $\mathbb{H}^3$ with different H values, which are asymptotic to the same asymptotic curve in $S_\infty^2(\mathbb{H}^3)$ are disjoint. In particular, for a given Jordan curve Γ in $S_\infty^2(\mathbb{H}^3)$, and given $-1 < H_1 < H_2 < 1$, it can be showed that the minimizing H -planes $\mathcal{P}_{H_1}$ and $\mathcal{P}_{H_2}$ in $\mathbb{H}^3$ with $\mathcal{P}_{H_i} = \Gamma$ are disjoint by using the ideas in [C4]. Hence, if Γ bounds a unique minimizing H -plane for any $H \in (-1, 1)$, it can be showed that the family of planes $\mathcal{F}_\Gamma = \{\mathcal{P}_H \mid \partial_\infty \mathcal{P}_H = \Gamma \text{ and } -1 < H < 1\}$ foliates $\mathbb{H}^3$. By the disjointness of the planes $\{\mathcal{P}_H\}$ for different H 's, in order to get the foliation, all one needs to show is that there is no gap between the planes $\{\mathcal{P}_H\}$ by using the uniqueness. By the arguments in [C4], a gap between the planes, say between $\{\mathcal{P}_H \mid -1 < H \leq H_0\}$ and $\{\mathcal{P}_H \mid H_0 < H < 1\}$, implies the nonuniqueness for H_0 -planes with $\mathcal{P}_{H_0} = \Gamma$. This gives a contradiction, and shows that $\mathcal{F}_\Gamma$ foliates $\mathbb{H}^3$. For example, if Γ is a star-shaped curve in $S_\infty^2(\mathbb{H}^3)$, then the family $\mathcal{F}_\Gamma$ foliates $\mathbb{H}^3$ [C4, GS, RT2].

REFERENCES

- [A1] M. Anderson, *Complete minimal varieties in $\mathbb{H}^n$* , Invent. Math. (1982) 477–494.
- [A2] M. Anderson, *Complete minimal hypersurfaces in hyperbolic n -manifolds*, Comment. Math. Helv. **58**, (1983) 264–290.
- [CMT] B. Coskunuzer, W. H. Meeks III, and G. Tinaglia, *Non-properly Embedded H -Planes in $\mathbb{H}^3$* , arXiv:1503.04641
- [C1] B. Coskunuzer, *Least Area Planes in Hyperbolic 3-Space are Properly Embedded*, Indiana Univ. Math. J. **58** (2009) 379–390.
- [C2] B. Coskunuzer, *Asymptotic Plateau Problem: A Survey*, Proc. Gokova GT Conf. (2013) 120–146.
- [C3] B. Coskunuzer, *Minimizing Constant Mean Curvature Hypersurfaces in Hyperbolic Space*, Geom. Dedicata **118** (2006) 157–171.
- [C4] B. Coskunuzer, *Foliations of hyperbolic space by constant mean curvature hypersurfaces*, IMRN (2010) 1417–1431.
- [C5] B. Coskunuzer, *Embeddedness of the Solutions of the H -Plateau Problem*, arXiv:1504.00661
- [Cu] T. Cuschiari, *Complete noncompact CMC surfaces in hyperbolic 3-space*, PhD thesis (2009) University of Warwick.
- [Ga] D. Gabai, *On the geometric and topological rigidity of hyperbolic 3-manifolds*, J. Amer. Math. Soc. **10** (1997) 37–74.
- [Gu] R. D. Gulliver, *The Plateau problem for surfaces of prescribed mean curvature in a Riemannian manifold*, J. Differential Geometry **8** (1973) 317–330.
- [GS] B. Guan, and J. Spruck, *Hypersurfaces of CMC in hyperbolic space with prescribed asymptotic boundary at infinity*, Amer. J. Math. **122**, (2000) 1039–1060.
- [MY] W.H. Meeks and S.T. Yau, *The classical Plateau problem and the topology of three-dimensional manifolds*, Topology **21** (1982) 409–442.

- [NS] B. Nelli and J. Spruck, *On the existence and uniqueness of CMC hypersurfaces in hyperbolic space*, Geometric analysis and the calculus of variations, 253–266, Internat. Press, Cambridge, MA, 1996.
- [RT1] J. Ripoll, F. Tomi, *Complete minimal discs in Hadamard manifolds*, arXiv:1409.0138
- [RT2] J. Ripoll, M. Telichevesky, *On the asymptotic Plateau problem for CMC hypersurfaces in hyperbolic space*, arXiv:1503.08083
- [To] Y. Tonegawa, *Existence and regularity of constant mean curvature hypersurfaces in hyperbolic space*, Math. Z. 221, (1996) 591–615.
- [Wa] B. Wang, *Least area Spherical Catenoids in $\mathbb{H}^3$* , arXiv:1204.4943

MIT, MATHEMATICS DEPARTMENT, CAMBRIDGE, MA 02139

KOC UNIVERSITY, DEPARTMENT OF MATHEMATICS, ISTANBUL 34450 TURKEY
E-mail address: bcoskunuzer@ku.edu.tr